



[cancer.org](https://www.cancer.org) | 1.800.227.2345

Treating Breast Cancer

If you've been diagnosed with breast cancer, your cancer care team will discuss your treatment options with you. It's important that you think carefully about each of your choices and weigh the benefits of each treatment option against the possible risks and side effects.

Local treatments

Some treatments, like surgery and radiation, are **local**, meaning they treat the tumor without affecting the rest of the body.

Most women with breast cancer will have some type of surgery to remove the tumor. Depending on the type of breast cancer and how advanced it is, you might need other types of treatment as well, either before or after surgery, or sometimes both.

- [Surgery for Breast Cancer](#)
- [Radiation for Breast Cancer](#)

Systemic treatments

Drugs used to treat breast cancer are considered **systemic therapies** because they can reach cancer cells almost anywhere in the body. Some can be given by mouth, injected into a muscle, or put directly into the bloodstream. Depending on the type of breast cancer, different types of drug treatment might be used, including:

- [Chemotherapy for Breast Cancer](#)
- [Hormone Therapy for Breast Cancer](#)
- [Targeted Drug Therapy for Breast Cancer](#)
- [Immunotherapy for Breast Cancer](#)

Common treatment approaches

Typically, treatment is based on the type of breast cancer and its stage. Other factors, including your overall health, menopause status, and personal preferences are also taken into account.

- [Treatment of Breast Cancer by Stage](#)
- [Treatment of Triple-negative Breast Cancer](#)
- [Treatment of Inflammatory Breast Cancer](#)
- [Treating Breast Cancer During Pregnancy](#)

Who treats breast cancer?

Based on your treatment options, you might have different types of doctors on your treatment team. These doctors could include:

- A **breast surgeon** or **surgical oncologist**: a doctor who uses surgery to treat breast cancer
- A **radiation oncologist**: a doctor who uses radiation to treat cancer
- A **medical oncologist**: A **surgeon**

you choose.

- [Questions to Ask Your Doctor About Breast Cancer](#)
- [Breast Reconstruction Surgery](#)
- [Seeking a Second Opinion](#)

Connect with a breast cancer survivor

[Reach To Recovery](#)

The American Cancer Society Reach To Recovery® program connects people facing breast cancer – from diagnosis through survivorship – with trained volunteers who are breast cancer survivors. Our volunteers provide one-on-one support through our website and mobile app to help those facing breast cancer cope with diagnosis, treatment, side effects, and more.

Thinking about taking part in a clinical trial

Clinical trials are carefully controlled research studies that are done to get a closer look at promising new treatments or procedures. Clinical trials are one way to get state-of-the-art cancer treatment. In some cases they may be the only way to get access to newer treatments. They are also the best way for doctors to learn better methods to treat cancer.

If you would like to learn more about clinical trials that might be right for you, start by asking your doctor if your clinic or hospital conducts clinical trials.

- [Clinical Trials](#)

Considering complementary and alternative methods

You may hear about alternative or complementary methods to relieve symptoms or treat your cancer that your doctors haven't mentioned. These methods can include vitamins, herbs, and special diets, or other methods such as acupuncture or massage, to name a few.

Complementary methods are treatments that are used **along with** your regular medical care. **Alternative** treatments are used **instead of** standard medical treatment. Although some of these methods might be helpful in relieving symptoms or helping you feel better, many have not been proven to work. Some might even be harmful.

Be sure to talk to your cancer care team about any method you are thinking about

using. They can help you learn what is known (or not known) about the method, which can help you make an informed decision.

- [Complementary and Integrative Medicine](#)

Help getting through cancer treatment

People with cancer need support and information, no matter what stage of illness they may be in. Knowing all of your options and finding the resources you need will help you make informed decisions about your care.

Whether you are thinking about treatment, getting treatment, or not being treated at all, you can still get supportive care to help with pain or other symptoms. Communicating with your cancer care team is important so you understand your diagnosis, what treatment is recommended, and ways to maintain or improve your quality of life.

Different types of programs and support services may be helpful, and they can be an important part of your care. These might include nursing or social work services, financial aid, nutritional advice, rehab, or spiritual help.

The American Cancer Society also has programs and services - including rides to treatment, lodging, and more - to help you get through treatment. Call our Cancer Knowledge Hub at 1-800-227-2345 and speak with one of our caring, trained cancer helpline specialists. Or, if you prefer, you can use our chat feature on cancer.org to connect with one of our specialists.

- [Palliative Care](#)
- [Programs & Services](#)

Choosing to stop treatment or choosing no treatment at all

For some people, when treatments have been tried and are no longer controlling the cancer, it could be time to weigh the benefits and risks of continuing to try new treatments. Whether or not you continue treatment, there are still things you can do to help maintain or improve your quality of life.

Some people, especially if the cancer is advanced, might not want to be treated at all. There are many reasons you might decide not to get cancer treatment, but it's important to talk to your doctors as you make that decision. Remember that even if you choose not to treat the cancer, you can still get supportive care to help with pain or other symptoms.

- [If Cancer Treatments Stop Working](#)

The treatment information given here is not official policy of the American Cancer Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor. Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask your cancer care team any questions you may have about your treatment options.

Surgery for Breast Cancer

Most women with breast cancer have some type of surgery as part of their treatment. There are different types of breast surgery, and they may be done for different reasons, depending on the situation. For example, surgery may be done to:

- Remove as much of the cancer as possible (breast-conserving surgery or mastectomy)
- Find out whether the cancer has spread to the lymph nodes under the arm (sentinel lymph node biopsy or axillary lymph node dissection)
- Restore the breast's shape after the cancer is removed (breast reconstruction)
- Relieve symptoms of advanced cancer

Your doctor may recommend a certain operation based on your breast cancer features and your medical history, or you may have a choice about which type of surgery to have. It's important to know your options so you can talk about them with your doctor and make the choice that is right for you.

- [Surgery to remove breast cancer](#)
- [Surgery to remove nearby lymph nodes](#)
- [Wire localization to guide surgery](#)
- [Breast reconstruction after surgery](#)
- [Surgery for advanced breast cancer](#)
- [More information about Surgery](#)

Surgery to remove breast cancer

More information about Surgery

For more general information about surgery as a treatment for cancer, see [Cancer Surgery](#)⁶.

<https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on June 25, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 4.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on June 25, 2021.

Last Revised: January 11, 2023

Breast-conserving Surgery (Lumpectomy)


Breast-conserving surgery (BCS) removes the cancer while leaving as much normal breast as possible. Usually, some surrounding healthy tissue and lymph nodes also are removed. Breast-conserving surgery is sometimes called **lumpectomy**, **quadrantectomy**, **partial mastectomy**, or **segmental mastectomy** depending on how much tissue is removed.

- [What you should know before having breast-conserving surgery](#)
- [Who can have breast-conserving surgery?](#)
- [Recovering from breast-conserving surgery](#)
- [Possible side effects of breast-conserving surgery](#)
- [Was all the cancer removed?](#)
- [Breast reconstruction surgery after breast-conserving surgery](#)
- [Treatment after breast-conserving surgery](#)

What you should know before having breast-conserving surgery

- How much of the breast is removed depends on the size and location of the tumor, your breast size, and other factors.
- Breast-conserving surgery allows a woman to keep most of her breast, but makes it likely she will also need [radiation](#).
- After BCS, most women will have radiation therapy. Some women might also get other treatments, such as [hormone therapy](#) or [chemotherapy](#).

- Choosing BCS plus radiation over mastectomy does not affect a woman's chances of long-term survival.
- If you think you want [breast reconstruction](#)¹, talk to your doctor **before** your breast cancer surgery.
- Not all women with breast cancer can have BCS. Talk to your doctor to find out whether BCS is an option for you.
- Side effects of BCS may include pain, a scar and/or dimple where the tumor was removed, a firm or hard surgical scar, and sometimes [lymphedema](#)², a type of swelling, in the arm.



Who can have breast-conserving surgery?

Breast-conserving surgery (BCS) is a good option for many women with early-stage

radiation oncologist for evaluation because each patient's cancer is unique.

BCS might be a good option if you:

- Are concerned about losing a breast
- Are willing to have radiation therapy and are able to get to the appointments (if you need help getting to and from your appointments see [Road To Recovery](#)³).
- Have not already had that breast treated with radiation therapy or BCS
- Have only one area of cancer in the breast, or multiple areas in one quadrant (multifocal) that are close enough to be removed together without changing the look of the breast too much
- Have a tumor smaller than 5 cm (2 inches), that is also small relative to the size of the breast
- Are not pregnant or, if pregnant, will not need radiation therapy immediately (to avoid risking harm to the fetus)
- Do not have a gene mutation (change) such as a *BRCA* or *ATM* mutation, which might increase your chance of a second breast cancer
- Do not have certain serious connective tissue diseases such as scleroderma or Sjögren's syndrome, which may make you very sensitive to the side effects of radiation therapy
- Do not have [inflammatory breast cancer](#)⁴
- Do not have positive margins (see *Was all the cancer removed?* below)

Recovering from breast-conserving surgery

This type of surgery is typically done in an outpatient surgery center, and an overnight stay in the hospital usually is not needed. Most women should be able to function after going home and can often return to their regular activities within 2 weeks. Some women may need help at home depending on how extensive their surgery was.

- Tips on bathing and showering after surgery
- When to call the doctor or nurse
- When to start using your arm again and how to do [arm exercises](#) to prevent stiffness
- When you can start wearing a bra again
- The use of medicines, including pain medicines and possibly antibiotics
- Any restrictions on activity
- What to expect regarding sensations or numbness in the breast and arm
- What to expect regarding feelings about body image
- When to see your doctor for a follow-up appointment
- Referral to a Reach To Recovery volunteer. Through our [Reach To Recovery program](#)⁵, a specially trained volunteer who has had breast cancer can provide information, comfort, and support.

Possible side effects of breast-conserving surgery

As with all operations, bleeding and infection at the surgery site are possible. Other side effects of breast-conserving surgery can include:

- Pain or tenderness or a "tugging" sensation in the breast
- Temporary swelling of the breast
- Hard scar tissue and/or a dimple that forms at the surgical site
- Swelling of the breast from a collection of fluid (seroma) that might need to be drained
- Change in the shape of the breast
- Neuropathic (nerve) pain (sometimes described as burning or shooting pain) in the chest wall, armpit, and/or arm that doesn't go away over time. This can also happen in mastectomy patients and is called [post-mastectomy pain syndrome](#)⁶ or PMPS.
- If [axillary lymph nodes are also removed](#), other side effects such as [lymphedema](#)⁷ may occur.

Was all the cancer removed?

During BCS, the surgeon will try to remove all the cancer, plus some surrounding normal tissue. This can sometimes be difficult depending on where the cancer is located in your breast.

After surgery, a doctor, called a pathologist, will look closely at the tissue that was removed in the lab. If the pathologist finds no invasive cancer cells at any of the edges of the removed tissue, it is said to have **negative** or clear margins. For women with DCIS, at least 2mm (0.08 inches) of normal tissue between the cancer and the edge of the removed tissue is needed. If the pathologist finds cancer cells at the edges of the removed tissue, it is said to have **positive** margins. This means that some cancer cells may still be in the breast after surgery, so the surgeon often needs to go back and remove more tissue. This operation is called a **re-excision**. If cancer cells are still found at the edges of the removed tissue after the second surgery, a mastectomy might be needed.

Having a **positive** margin means that some cancer cells may still be in the breast after surgery, so the surgeon often needs to go back and remove more tissue. This operation is called a **re-excision**. If cancer cells are still found at the edges of the removed tissue after the second surgery, a mastectomy might be needed.

Breast reconstruction surgery after breast-conserving surgery

OJ Vilholm, S Cold, L Rasmussen and SH Sindrup. The postmastectomy pain syndrome: an epidemiological study on the prevalence of chronic pain after surgery for breast cancer. *British Journal of Cancer* (2008) 99, 604 – 610.

Sabel MS. Breast-conserving therapy. In Chen W, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Accessed July 7, 2021.

Last Revised: October 27, 2021

Mastectomy

Mastectomy is breast cancer surgery that removes the entire breast.

A mastectomy might be done:

- When a woman cannot be treated with [breast-conserving surgery \(lumpectomy\)](#), which saves most of the breast.
 - If a woman chooses mastectomy over breast-conserving surgery for personal reasons.
 - For women at very high risk of getting a second breast cancer who sometimes choose to have a double mastectomy (the removal of both breasts).
-
- [Types of mastectomies](#)
 - [Who might get a mastectomy?](#)
 - [Breast reconstruction surgery after mastectomy](#)
 - [Going flat](#)
 - [Recovering from a mastectomy](#)
 - [Possible side effects of mastectomy](#)
 - [Treatment after mastectomy](#)

Types of mastectomies


There are several different types of mastectomies, based on how the surgery is done and how much tissue is removed.

Simple (or total) mastectomy

In this procedure, the surgeon removes the entire breast, including the nipple, areola, fascia (covering) of the pectoralis major muscle (main chest muscle), and skin. A few underarm lymph nodes might be removed as part of a [sentinel lymph node biopsy](#) depending on the situation. Most women, if they are hospitalized, can go home the next day.

Modified radical mastectomy

A modified radical mastectomy combines a simple mastectomy with the removal of the lymph nodes under the arm (called an [axillary lymph node dissection](#)).



Radical mastectomy

This extensive surgery is rarely done now. The surgeon removes the entire breast, axillary (underarm) lymph nodes, and the pectoral(chest wall) muscles under the breast. This surgery was once very common, but less extensive surgery (such as the modified

radical mastectomy) has been found to be just as effective and with fewer side effects. This operation might be done if the tumor is growing into the pectoral muscles.

Skin-sparing mastectomy

In this procedure, most of the skin over the breast is left place. Only the breast tissue, nipple, and areola are removed. The amount of breast tissue removed is the same as with a simple mastectomy. Implants or tissue from other parts of the body can be used during the surgery to [reconstruct the breast](#)¹.

Many women prefer a skin-sparing mastectomy because it offers the advantage of less scar tissue and a reconstructed breast that seems more natural. But it may not be suitable for larger tumors or those that are close to the surface of the skin.

The risk of local cancer recurrence with this type of mastectomy is the same as with other types of mastectomies.

Experts recommended that skin-sparing mastectomies be done by a team of breast surgeons with a lot of experience in this procedure.

Nipple-sparing mastectomy

A nipple-sparing mastectomy is similar to a skin-sparing mastectomy in that the breast tissue is removed and the breast skin is saved. But in this procedure, the nipple and areola are left in place. This can be followed by breast reconstruction. The surgeon often removes the breast tissue under the nipple and areola during the procedure to check for cancer cells. If cancer is found in this tissue, the nipple and areola must be removed.

This type of mastectomy is more often an option for women who have a small, early-stage cancer, away (more than 2cm) from the nipple and areola, with no signs of cancer in the skin or the nipple.

As with any surgery, there are risks. After the surgery, the nipple may not have a good blood supply, causing the tissue to shrink or become deformed. Because the nerves are also cut, there often may be little or no feeling left in the nipple. If a woman has larger breasts, the nipple may look out of place after the breast is reconstructed. As a result, many doctors feel that this surgery is best done in women with small to medium sized breasts. This procedure leaves fewer scars you can see, but it also has a risk of leaving behind more breast tissue than other forms of mastectomy. This could result in a higher risk of cancer developing than for a skin-sparing or simple mastectomy. However,

improvements in technique have helped lower this risk and the risk of cancer coming back in the same area is about the same as with other types of mastectomies. Most experts consider nipple-sparing mastectomy to be an acceptable treatment for breast cancer in certain cases.

As with a skin-sparing mastectomy, experts also recommended that this type of mastectomy be done by a team of breast surgeons with a lot of experience with this procedure.

Double mastectomy

When both breasts are removed, it is called a **double (or bilateral) mastectomy**. Double mastectomy is sometimes done as a risk-reducing (or preventive) surgery for women at very high risk for getting breast cancer, such as those with a *BRCA* gene mutation. Most of these mastectomies are simple mastectomies, but some may be nipple-sparing. There are other situations where a double mastectomy might be done as part of a woman's breast cancer treatment plan. This is done after careful consideration and discussion between the patient and their cancer care team.

Who might get a mastectomy?

Many women with early-stage cancers can choose between breast-conserving surgery (BCS) and mastectomy. You may prefer mastectomy as a way to "take out all the cancer as quickly as possible." But the fact is that in most cases, mastectomy does not give you any better chance of long-term survival compared to BCS. Studies of thousands of women over more than 20 years show that when BCS is done along with radiation, the outcome is the same as having a mastectomy.

Mastectomy might be recommended if you:

- Are unable to have radiation therapy
- Would prefer more extensive surgery instead of having radiation therapy
- Have had the breast treated with radiation therapy in the past
- Have already had BCS with re-excision(s) that did not completely remove the cancer
- Have two or more areas of cancer in different quadrants of the same breast (multicentric) that are not close enough to be removed together without changing the look of the breast too much
- Have a tumor larger than 5 cm (2 inches) across, or a tumor that is large relative to your breast size

- Are pregnant and would need radiation therapy while still pregnant (risking harm to the fetus)
- Have a genetic factor such as a *BRCA* mutation, which might increase your chance of a second cancer
- Have a serious connective tissue disease such as scleroderma or lupus, which may make you especially sensitive to the side effects of radiation therapy
- Have inflammatory breast cancer

For women who are worried about breast cancer coming back, it is important to understand that having a mastectomy instead of breast-conserving surgery plus radiation **only** lowers your risk of developing a second breast cancer in the same breast. It does not lower the chance of the cancer coming back in other parts of the body, including the opposite breast.

Breast reconstruction surgery after mastectomy

After having a mastectomy a woman might want to consider having the breast mound rebuilt to restore the breast's appearance. This is called [breast reconstruction](#)². Although each case is different, most mastectomy patients can have reconstruction. Reconstruction can be done at the same time as the mastectomy or sometime later.

If you are thinking about having reconstructive surgery, it's a good idea to discuss it with your surgeon and a plastic surgeon before your mastectomy. This allows the surgical teams to plan the treatment that's best for you, even if you wait and have the reconstructive surgery later. Insurance companies typically cover breast reconstruction, but you should check with your insurance company so you know what is covered.

Going flat

Some women [choose not to have reconstructive surgery](#)³. Wearing a breast prosthesis (breast form) is an option for women who want to have the shape of a breast under their clothes without having surgery. Some women are also comfortable with just '[going flat](#)⁴'. Going flat involves a procedure called **aesthetic flat closure** or **flat closure**. A flat closure means the extra fat, skin, and other tissue in the breast area are removed and the leftover tissue is tightened and smoothed out to flatten the chest wall.

Recovering from a mastectomy

In general, women having a mastectomy stay in the hospital for 1 or 2 nights and then go home. How long it takes to recover from surgery depends on what procedures were done, and some women may need help at home. Most women should be fairly functional after going home and can often return to their regular activities within about 4 weeks. Recovery time is longer if breast reconstruction was also done, and it can take months to return to full activity after some procedures.

Ask your health care team how to care for your surgery site and arm. Usually, you and your caregivers will get written instructions about care after surgery. These instructions typically cover:

- How to care for the surgery site and dressing
- How to care for your drain, if you have one (this is a plastic or rubber tube coming out of the surgery site attached to a soft rubber ball that collects the fluid that occurs during healing)
- How to tell if an infection is starting
- Bathing and showering after surgery
- When to call the doctor or nurse
- When to start using your arm again and how to do [arm exercises](#) to prevent stiffness
- When you can start wearing a bra again
- When to begin using a prosthesis and what type to use
- Use of medicines, including pain medicines and possibly antibiotics
- Any restrictions on activity
- What to expect regarding sensations or numbness in the breast and arm
- What to expect regarding feelings about body image
- When to see your doctor for a follow-up appointment
- Referral to a Reach To Recovery volunteer. Through our [Reach To Recovery program](#)⁵, a specially trained volunteer who has had breast cancer and can provide information, comfort, and support.

Possible side effects of mastectomy

Bleeding and infection at the surgery site are possible with all operations. The side effects of mastectomy can depend on the type of mastectomy you have (complex surgeries tend to have more side effects). Side effects can include:

- Pain or tenderness at the surgery site

- Swelling at the surgery site
- Buildup of blood in the wound (hematoma)
- Buildup of clear fluid in the wound (seroma)
- Limited arm or shoulder movement
- Numbness in the chest or upper arm
- Neuropathic (nerve) pain (sometimes described as burning or shooting pain) in the chest wall, armpit, and/or arm that doesn't go away over time. It is also called [post-mastectomy pain syndrome or PMPS](#)⁶.
- If axillary lymph nodes are also removed, other side effects such as [lymphedema](#)⁷ may occur.

Treatment after mastectomy

Some women might get other treatments after a mastectomy, such as [hormone therapy](#) to help lower the risk of the cancer coming back. Some women might also need [chemotherapy](#), or [targeted therapy](#) after surgery. If so, [radiation therapy](#) and/or hormone therapy is usually delayed until the chemotherapy is completed. Talk to your doctor about what to expect.

Hyperlinks

1. www.cancer.org/cancer/types/breast-cancer/reconstruction-surgery/breast-reconstruction-options.html
2. www.cancer.org/cancer/types/breast-cancer/reconstruction-surgery.html
3. www.cancer.org/cancer/types/breast-cancer/reconstruction-surgery/breast-reconstruction-alternatives.html
4. www.cancer.org/cancer/types/breast-cancer/reconstruction-surgery/breast-reconstruction-alternatives.html
5. www.cancer.org/support-programs-and-services/reach-to-recovery.html
6. www.cancer.org/cancer/managing-cancer/side-effects/pain/post-mastectomy-pain-syndrome.html
7. www.cancer.org/cancer/managing-cancer/side-effects/swelling/lymphedema.html

References

Gieni M, Avram R, Dickson L, et al. Local breast cancer recurrence after mastectomy

and immediate breast reconstruction for invasive cancer: a meta-analysis. *Breast* 2012;21(3):230–236.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Kwong A and Sabel MS. Mastectomy. In Chen W, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Accessed July 8, 2021.

National Cancer Institute. Aesthetic flat closure. Accessed at <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/aesthetic-flat-closure> on December 20, 2023.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Patient Version. 2021. Accessed at <https://www.cancer.gov/types/breast/patient/breast-treatment-pdq> on July 8, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 4.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on June 25, 2021.


Oh J.L. (2008) Multifocal or Multicentric Breast Cancer: Understanding Its Impact on Management and Treatment Outcomes. In: Hayat M.A. (eds) *Methods of Cancer Diagnosis, Therapy and Prognosis*. *Methods of Cancer Diagnosis, Therapy and Prognosis*, vol 1. Springer, Dordrecht. <https://doi.org/10.1007/978-1-4020m> Elsevier; 2020.

Last Revised: December 20, 2023

Lymph Node Surgery for Breast Cancer

If breast cancer spreads, it typically goes first to nearby [lymph nodes](#)¹ under the arm. It can also sometimes spread to lymph nodes near the collarbone or near the breast bone (the front center of the chest). Knowing if the cancer has spread to your lymph nodes helps doctors find the best way to treat your cancer.

If you have been diagnosed with breast cancer, it's important to find out [how far the cancer has spread](#)². To help find out if the cancer has spread outside the breast, one or more of the lymph nodes under the arm (axillary lymph nodes) are removed and checked in the lab.



other lymph nodes in the same area will also have cancer if cancer is found in the sentinel lymph node(s), the surgeon may go ahead with an axillary dissection (ALND) to remove more lymph nodes while you are still on the operating table. If no cancer cells are seen in the node(s) at the time of the surgery, or if they are not checked by a pathologist at the time of the surgery, they will be examined more closely over the next several days.

If cancer is found in the sentinel node(s) later, the surgeon may recommend an ALND at a later time to check more nodes for cancer. Studies have shown, however, that in some cases it may be safe to leave the rest of the lymph nodes behind. This is based on certain factors, such as the size of the breast tumor, what type of surgery is used to remove the tumor, and what treatment is planned after surgery, among other things.

Based on the studies that have looked at this, skipping the ALND may be an option for:

- Women with breast tumors 5 cm (about 2 inches) across or smaller who have no more than 2 positive sentinel lymph nodes, are having breast-conserving surgery followed by radiation, and did not get any chemotherapy before surgery.
- Women who have lymph nodes with a very small amount of cancer (no more than 2 mm) and are having a [mastectomy](#).

If there is no cancer in the sentinel node(s), it's very unlikely that the cancer has spread to other lymph nodes, so no further lymph node surgery will be needed.

SLNB is often considered for women with early-stage breast cancer and is typically not used for women with inflammatory breast cancer. It might be used for women with locally advanced breast cancer in certain instances, such as after neoadjuvant treatment.

Although SLNB has become a common procedure, it requires a great deal of skill. It should be done only by a surgeon who has experience with this technique. If you are offered this type of biopsy, ask your surgeon if they do them regularly.


Axillary lymph node dissection (ALND)

In this procedure, anywhere from about 10 to 40 (though usually less than 20) lymph nodes are removed from the area under the arm (axilla) and checked for cancer spread. ALND is usually done at the same time as a mastectomy or [breast-conserving surgery \(BCS\)](#), but it can be done in a second operation. ALND may be needed:


- If a previous SLNB has shown 3 or more of the underarm lymph nodes have cancer

cells

- If swollen underarm or collarbone lymph nodes can be felt before surgery or seen on imaging tests and a FNA or core needle biopsy shows cancer
- If the cancer has grown large enough to extend outside the lymph node(s)
- If the SLNB is positive for cancer cells after chemotherapy was given to shrink the tumor before surgery



Side effects of lymph node surgery



Lymphedema is less common after a sentinel lymph node biopsy (SLNB) than an axillary lymph node dissection (ALND). The risk is thought to be in the range of 5% to 17% in women who have a SLNB and around 20% to 30% in women who have an ALND. It may be more common if radiation is given after surgery or in women who are obese. Sometimes the swelling lasts for only a few weeks and then goes away. But in some women, it lasts a long time. If your arm is swollen, tight, or painful after lymph node surgery, be sure to tell someone on your cancer care team right away.

Limited arm and shoulder movement

You might also have **limited movement in your arm and shoulder** after surgery. This is more common after ALND than SLNB. Your doctor may advise [exercises](#) to help keep you from having long-lasting problems (a frozen shoulder).

Some women notice a rope-like structure that begins under the arm and can extend down toward the elbow. This is sometimes called **axillary web syndrome** or **lymphatic**

cording. It is more common after ALND than SLNB. Symptoms may not appear for weeks or even months after surgery. It can cause pain and limit movement of the arm and shoulder. This often goes away without treatment, although some women may find physical therapy helpful.

Numbness

Numbness of the skin on the upper, inner arm is a common side effect because the nerve that controls sensation (feeling) here travels through the lymph node area.

Hyperlinks

1. www.cancer.org/cancer/diagnosis-staging/lymph-nodes-and-cancer.html
-

Biopsy in Breast Cancer Surgery. *Ann Surg Oncol*. 2016;23(11):3459–3466.
doi:10.1245/s10434-016-5432-8.

Exercises After Breast Cancer Surgery

- Mastectomy
- Breast reconstruction

Any of these can affect how well you can move your shoulder and arm or go about your daily activities, like dressing, bathing, and combing your hair. Pain and stiffness can cause weakness and limit movement of your arm and shoulder.

- [Exercises can help get movement back](#)
- [The week after surgery](#)
- [General guidelines for these exercises](#)
- [Things to keep in mind after breast surgery](#)
- [Other kinds of exercise](#)

Exercises can help get movement back

No matter what type of surgery you have, it's important to do exercises afterward to get the arm and shoulder moving again. Exercises help decrease side effects of your surgery and help you get back to your usual activities.

If you've had radiation therapy after surgery, exercises are even more important to help keep your arm and shoulder flexible. Radiation may affect your arm and shoulder long after treatment is finished. Because of this, it's important to develop a regular habit of doing exercises to maintain arm and shoulder mobility after radiation treatments for breast cancer.

It's very important to talk with your doctor before starting any exercises so that you can decide on a program that's right for you. Your doctor might suggest you see a physical therapist or occupational therapist, or a cancer exercise specialist certified by the American College of Sports Medicine. These health professionals are specially trained to design an exercise program just for you. You might need this kind of help if you do not have full use of your arm within 3 to 4 weeks of surgery.

Some exercises should not be done until drains and sutures (stitches) are removed, but others can be done soon after surgery. The exercises that increase your shoulder and arm motion can usually be started in a few days. Exercises to help make your arm stronger are added later.

Here are some of the more common exercises that women do after breast surgery. Talk to your doctor or therapist about which of these are right for you and when you should start doing them. Do not start any of these exercises without talking to your doctor first.

The week after surgery


The tips and exercises listed below should be done for the first 3 to 7 days after surgery. **Do not do them until you get the OK from your doctor.**

- Use your affected arm (on the same side as your surgery) as you normally would when you comb your hair, bathe, get dressed, and eat. Be sure to ask your doctor if you can lift heavier items.
- Raise your arms up to shoulder height, or whatever height you can without pulling on your drains, and lower them again a few times. Repeat this 3 or 4 times a day. These exercises help move lymph fluid out of your arm and help restore movement. You can exercise your affected arm while it's raised. There are a few ways to do this. With your arm raised, open and close your hand 15 to 25 times. Next, 7cf 1 95.35 517.02 Tm

- It may be helpful to exercise after a warm shower when muscles are warm and relaxed.
- Wear comfortable, loose clothing when doing the exercises.
- Do the exercises slowly until you feel a gentle stretch. Hold each stretch at the end of the motion and slowly count to 5. It's normal to feel some pulling as you stretch the skin that has been shortened because of the surgery. Do not bounce or make any jerky movements when doing any of the exercises. You should not feel pain as you do them, only gentle stretching.
- Do each exercise 5 to 7 times. Try to do each exercise correctly. Tell your cancer care team if you have trouble doing them . You may need to be referred to a physical or occupational therapist.
- Do the exercises twice a day until you get back your normal flexibility. Continuing to do some exercises during the months after surgery can help you keep moving.
- Be sure to take deep breaths, in and out, as you do each exercise.
- The exercises are set up so that you start them first lying down, then sitting, and finish them standing up.


Here are some of the more common exercises that women do after breast surgery. Talk to your doctor or therapist about which of these are right for you and when you should start doing them. Do not start any of these exercises without talking to your doctor first.

Wand exercise



This exercise helps increase your ability to move your shoulders forward. You will need a broom handle, yardstick, or other stick-like object to use as the wand in this exercise. Do these exercises on a bed or the floor. Lie on your back with your knees bent and your feet flat.


- Hold the wand across your belly in both hands with your palms facing up. Lift the wand up over your head as far as you can. Use your unaifo rg /GI tharmse a inc87.6 259.9



This exercise helps increase the movement in the front of your chest and shoulder. It may take many weeks of regular exercise before your elbows will get close to the bed or floor. Do these exercises on a bed or the floor. Lie on your back with your knees bent and your feet flat.

- Clasp your hands behind your neck with your elbows pointing toward the ceiling.
- Move your elbows apart and down toward the bed or floor.
- Repeat 5 to 7 times.


Shoulder blade stretch



This exercise helps increase your shoulder blade movement.

- Sit in a chair very close to a table with your back against the back of the chair.
- Place the unaffected arm on the table with your elbow bent and palm down. Do not move this arm during the exercise.
- Place the affected arm on the table, palm down, with your elbow straight.
- Without moving your trunk, slide the affected arm forward, toward the opposite side of the table. You should feel your shoulder blade move as you do this.
- Relax your arm and repeat 5 to 7 times.


Shoulder blade squeeze



This exercise also helps increase shoulder blade movement and improve posture.

- Sit in a chair in front of a mirror. Face straight ahead. Do not rest against the back of the chair.
- Your arms should be at your sides with your elbows bent.
- Squeeze your shoulder blades together, bringing your elbows behind you toward your spine. Elbows will move with you, but don't force the motion with your elbows. Keep your shoulders level as you do this. Do not lift your shoulders up toward your ears.
- Return to the starting position and repeat 5 to 7 times.


Side bends



This exercise helps increase movement of your trunk and body.

- Sit in a chair and clasp your hands together in front of you. Lift your arms slowly over your head, straightening your arms.
- When your arms are over your head, bend your trunk to the right keeping your arms overhead.
- Return to the starting position and bend to the left.
- Repeat 5 to 7 times.

Chest wall stretch



This exercise helps increase your mobility in your shoulder.

- Stand facing the wall with your toes about 8 to 10 inches from the wall.
- Put your hands on the wall. Use your fingers to "climb the wall," reaching as high as you can until you feel a stretch.
- Return to the starting position and repeat 5 to 7 times.
- The picture shows both arms going up at the same time, but you might find it easier to raise one arm at a time.
- Be sure you keep your shoulders dropped far away from your ears as you raise your arms. Keep your ears over your shoulders to avoid making your neck sore.

Things to keep in mind after breast surgery

Start exercising slowly and do more as you are able. Stop exercising and talk to your doctor right away if you:

- Get weaker, start losing your balance, or start falling
- Have pain that gets worse
- Have new heaviness, aching, tightness, or other strange sensations in your arm
- Have unusual swelling or swelling that gets worse
- Have headaches, dizziness, blurred vision, new numbness, or tingling in your arms or chest

McNeely ML, Campbell K, Ospina M, et al. Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database Syst Rev*. 2010;(6):CD005211.

Exercise and Lymphedema: Position Statement of the National Lymphedema Network. March 2018. Accessed at <https://www.mylymph.com/2018/03/08/exercise-and-lymphedema-position-statement-of-the-national-lymphedema-network> on August 12, 2019.

Petito EL, Nazário AC, Martinelli SE, Facina G, De Gutiérrez MG. Application of a domicile-based exercise program for shoulder rehabilitation after breast cancer surgery. *Rev Lat Am Enfermagem*. 2012;20(1):35-43.

Sagen A, Kaaresen R, Sandvik L, Thune I, Risberg MA. Upper limb physical function and adverse effects after breast cancer surgery: a prospective 2.5-year follow-up study and preoperative measures. *Arch Phys Med Rehabil*. 2014;95(5):875-881.

Last Revised: October 27, 2021

Radiation for Breast Cancer

Radiation therapy is treatment with high-energy rays (or particles) that destroy cancer cells. Some women with breast cancer will need radiation, in addition to other treatments.

Depending on the breast cancer's [stage](#)¹ and other factors, radiation therapy can be used in several situations:

- After breast-conserving surgery (BCS), to help lower the chance that the cancer will come back in the same breast or nearby lymph nodes.
- After a mastectomy, especially if the cancer was larger than 5 cm (about 2 inches), if cancer is found in many lymph nodes, or if certain surgical margins, such as the skin or muscle, have cancer cells.
- If cancer has spread to other parts of the body, such as the bones, spinal cord, or brain.
- [Types of radiation therapy for breast cancer](#)

- [External beam radiation therapy \(EBRT\)](#)
- [Types and schedules of external beam radiation for breast cancer](#)
- [Brachytherapy](#)
- [Types of brachytherapy](#)
- [More information about radiation therapy](#)

Types of radiation therapy for breast cancer

The main types of radiation therapy that can be used to treat breast cancer are:

- External beam radiation therapy
- Brachytherapy

External beam radiation therapy (EBRT)

[EBRT²](#) is the most common type of radiation therapy for women with breast cancer. A machine outside the body focuses the radiation on the area affected by the cancer.

Which areas need radiation depends on whether you had a [mastectomy](#) or [breast-conserving surgery](#) (BCS) and if the cancer has reached nearby lymph nodes.

- If you had a mastectomy and no lymph nodes had cancer cells, radiation will be focused on the chest wall, the mastectomy scar, and the places where any drains exited the body after surgery.
If you had BCS, you will most likely have radiation to the entire breast (called **whole breast radiation**). An extra **boost** of radiation to the area in the breast where the cancer was removed (called the **tumor bed**) is often given if there is a high risk of the cancer coming back. The boost is often given after the treatments to the whole breast have ended. It uses the same machine, with lower amounts of radiation

- **3D-conformal radiotherapy (3D-CRT):** In this technique, the radiation is given with special machines so that it is better aimed at the tumor bed. This spares more of the surrounding normal breast tissue. Treatments are given twice a day for 5 days or daily for 2 weeks.
- **Intensity-modulated radiotherapy (IMRT):** IMRT is like 3D-CRT, but it also changes the strength of some of the beams in certain areas. This gets stronger doses to certain parts of the tumor bed and helps lessen damage to nearby normal body tissues.
- **Brachytherapy:** See brachytherapy below.

Women who are interested in these approaches may want to ask their doctor about taking part in [clinical trials](#)³ of accelerated partial breast irradiation.

Chest wall radiation

If you had a mastectomy and none of the lymph nodes had cancer, radiation will be given to the entire chest wall, the mastectomy scar, and the areas of any surgical drains. It is typically given every day, 5 days a week, for 6 weeks.

Lymph node radiation

Whether or not you have had BCS or a mastectomy, if cancer was found in the lymph

because it could make the skin changes worse. Most skin changes get better within a few months. Changes to the breast tissue usually go away in 6 to 12 months, but it can take longer.

External beam radiation therapy can also cause side effects later on:

- Some women may find that radiation therapy causes the breast to become smaller and the skin to become firmer or swollen.
- Radiation may affect your options for breast reconstruction later on. It can also raise the risk of problems with appearance and healing if it's given after reconstruction, especially tissue flap procedures.
- Women who have had breast radiation may not be able to breastfeed from the radiated breast.
- Radiation to the breast can sometimes damage some of the nerves to the arm. This is called **brachial plexopathy** and can lead to numbness, pain, and weakness in the shoulder, arm, and hand.
- Radiation to the underarm lymph nodes might cause [lymphedema](#)⁴, a type of pain and swelling in the arm or chest.
- In rare cases, radiation therapy may weaken the ribs, which could lead to a fracture.
- In the past, parts of the lungs and heart were more likely to get some radiation, which could lead to long-term damage of these organs in some women. Modern radiation therapy equipment focuses the radiation beams better than older machines, so these problems are rare today.
- A very rare complication of radiation to the breast is the development of another cancer called an [angiosarcoma](#)⁵.

Brachytherapy

[Brachytherapy](#)⁶, also known as **internal radiation**, is another way to deliver radiation therapy. Instead of aiming radiation beams from outside the body, a device containing

Types of brachytherapy

Intracavitary brachytherapy

Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 7.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 30, 2021.

Shah C, Vicini F, Shaitelman S, Hepel J, Keisch M, Arthur D et al. The American Brachytherapy Society consensus statement for accelerated partial-breast irradiation. *Brachytherapy*.2018; 17(1), 154–170.

Shaitelman SF, Schlembach PJ, Arzu I, et al. Acute and short-term toxic effects of conventionally fractionated vs hypofractionated whole-breast irradiation: A randomized clinical trial. *JAMA Oncol*. 2015;1:931-941.

Smith GL, Xu Y, Buchholz TA, et al. Association between treatment with brachytherapy vs whole-breast irradiation and subsequent mastectomy, complications, and survival among older women with invasive breast cancer. *JAMA*. 2012;307:1827-1837.

Stmad V, Ott OJ, Hildebrandt G, et al. 5-year results of accelerated partial breast irradiation using sole interstitial multicatheter brachytherapy versus whole-breast irradiation with boost after breast-conserving surgery for low-risk invasive and in-situ carcinoma of the female breast: a randomised, phase 3, non-inferiority trial. *Lancet*. 2016 Jan 16;387(10015):229-38.

Taghian A. Adjuvant radiation therapy for women with newly diagnosed, non-metastatic breast cancer. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated May 7, 2021. Accessed August 30, 2021.

Taghian A. Radiation therapy techniques for newly diagnosed, non-metastatic breast cancer. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated November 12, 2021. Accessed August 30, 2021.

Whelan TJ, Pignol J, Levine MN, et al. Long-Term Results of Hypofractionated Radiation Therapy for Breast Cancer. *N Engl J Med* 2010; 362:513-520.

Chemotherapy for Breast Cancer

cancers that are too big to be removed by surgery when first diagnosed, have many lymph nodes involved with cancer, or are inflammatory breast cancers .

If after neoadjuvant chemo, cancer cells are still found when surgery is done (also called residual disease), you might be offered more chemotherapy (adjuvant chemotherapy) to reduce the chances of the cancer coming back (recurrence).

Some other reasons you might get neoadjuvant chemo:

- By giving chemo before the tumor is removed, doctors can see how the cancer responds to it. If the first set of chemo drugs doesn't shrink the tumor, your doctor will know that other drugs are needed. It should also kill any cancer cells that might have spread but can't be seen by the naked eye or on imaging tests. Just like adjuvant chemo, neoadjuvant chemo can lower the risk of breast cancer coming back.
- Some people with early-stage cancer who get neoadjuvant chemo might live longer if the cancer completely goes away with that treatment. This can be seen most often in women who have triple-negative breast cancer or HER2-positive breast cancer.
- Getting chemo before surgery can also give some people extra time to get genetic testing or plan reconstructive surgery.

Keep in mind that not all women with breast cancer are good candidates for neoadjuvant chemo.

For metastatic breast cancer

Chemo can be used as the main treatment for women whose cancer has spread outside the breast and underarm area to distant organs like the liver or lungs. Chemo can be given either when breast cancer is diagnosed or after initial treatments. The length of treatment depends on how well the chemo is working and how well you tolerate it.

Chemotherapy drugs used for breast cancer

In most cases, chemo has the greatest effect when more than one drug is used at a time. Often, combinations of 2 or 3 drugs are used. Doctors use many different combinations, and it's not clear that any particular drug combination is the best.

Adjuvant and neoadjuvant chemo drugs

- Anthracyclines, such as doxorubicin (Adriamycin) and epirubicin (Ellence)
- Taxanes, such as paclitaxel (Taxol) and docetaxel (Taxotere)
- 5-fluorouracil (5-FU) or capecitabine (Xeloda)
- Cyclophosphamide (Cytoxan)
- Carboplatin (Paraplatin)

Chemo drugs for breast cancer that has spread (metastatic breast cancer)

- Taxanes: Paclitaxel (Taxol), docetaxel (Taxotere), and albumin-bound paclitaxel (Abraxane)
- Ixabepilone (Ixempra)
- Eribulin (Halaven)
- Anthracyclines: Doxorubicin (Adriamycin), liposomal doxorubicin (Doxil), and epirubicin (Ellence)
- Platinum agents (Cisplatin, carboplatin)
- Vinorelbine (Navelbine)
- Capecitabine (Xeloda)
- Gemcitabine (Gemzar)
- [Antibody drug conjugates](#) (Ado-trastuzumab emtansine [Kadcyla], Fam-trastuzumab deruxtecan [Enhertu], Sacituzumab govitecan [Trodelvy])

Although drug combinations are often used to treat early breast cancer, advanced

chemo. These are known as [central venous catheters](#)⁴ (CVCs), central venous access devices (CVADs), or central lines. They are used to put medicines, blood products, nutrients, or fluids right into your blood. They can also be used to take out blood for testing.

There are many different kinds of CVCs. The most common types are the port and the PICC line. For breast cancer patients, the central line is typically placed on the side opposite of the breast cancer. If a woman has breast cancer in both breasts, the central

If you think you might want to have children after being treated for breast cancer, talk with your doctor soon after being diagnosed and before you start treatment. For some women, adding medicines, like monthly injections with a [luteinizing hormone-releasing hormone \(LHRH\) analog](#), along with chemo, can help them have a successful pregnancy after cancer treatment. To learn more, see [Female Fertility and Cancer](#)⁷

syndrome. Early symptoms include numbness, tingling, and redness. If it gets worse, the hands and feet can become swollen and uncomfortable or even painful. The skin may blister, leading to peeling or even open sores. There is no specific treatment, although some creams or steroids given before chemo may help. These symptoms gradually get better when the drug is stopped or the dose is lowered. The best way to prevent severe hand-foot syndrome is to tell your doctor when symptoms first come up,

see [Chemotherapy](#)¹⁰.

To learn about some of the side effects listed here and how to manage them, see [Managing Cancer-related Side Effects](#)¹¹.

Hyperlinks

Greer AC, Lanes A, Poorvu PD, et al. The impact of fertility preservation on the timing of breast cancer treatment, recurrence, and survival [published online ahead of print, 2021 Jun 23]. *Cancer*. 2021;10.1002/cncr.33601. doi:10.1002/cncr.33601.

Hermelink K. Chemotherapy and Cognitive Function in Breast Cancer Patients: The So-Called Chemo Brain. *J Natl Cancer Inst Monogr*. 2015 May;2015(51):67-9.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Health Professional Version. 2021. Accessed at <https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on August 10, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 6.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 10, 2021.

Osborne CK. Chapter 53: Adjuvant Systemic Therapy Treatment Guidelines. In: Harris JR, Lippman ME, Morrow M, Osborne CK, eds. *Diseases of the Breast*. 5th ed. Philadelphia: Wolters Kluwer Health; 2014.

Last Revised: October 27, 2021

Hormone Therapy for Breast Cancer

Some types of breast cancer are affected by hormones, like estrogen and progesterone. The breast cancer cells have receptors (proteins) that attach to estrogen and progesterone, which helps them grow. Treatments that stop these hormones from attaching to these receptors are called **hormone or endocrine therapy**.

Hormone therapy can reach cancer cells almost anywhere in the body and not just in

the breast. It's recommended for women with tumors that are hormone receptor-positive. It does not help women whose tumors don't have hormone receptors (these tumors are called hormone receptor-negative).

- [When is hormone therapy used for breast cancer?](#)
- [How does hormone therapy work?](#)
- [Drugs that block estrogen receptors](#)
- [Drugs that lower estrogen levels](#)
- [Hormone therapy after surgery for breast cancer](#)
- [If cancer comes back or has spread](#)
- [Less common types of hormone therapy](#)
- [More information about hormone therapy](#)

When is hormone therapy used for breast cancer?

Hormone therapy is often used after surgery (as adjuvant therapy) to help reduce the risk of the cancer coming back. Sometimes it is started before surgery (as neoadjuvant therapy).

It is usually taken for at least 5 years. Treatment longer than 5 years might be offered to women whose cancers have a higher chance of coming back. A test called the [Breast Cancer Index](#)¹ might be used to help decide if a woman will benefit from more than 5 years of hormone therapy.

Hormone therapy can also be used to treat cancer that has come back after treatment or that has spread to other parts of the body.

How does hormone therapy work?

About 2 out of 3 breast cancers are hormone receptor-positive. Their cells have receptors (proteins) for estrogen (ER-positive cancers) and/or progesterone (PR-positive cancers) which help the cancer cells grow and spread.

There are several types of hormone therapy for breast cancer. Most types of hormone therapy either lower estrogen levels in the body or stop estrogen from helping breast cancer cells grow.

Drugs that block estrogen receptors

These drugs work by stopping estrogen from fueling breast cancer cells to grow.

Selective estrogen receptor modulators (SERMs)

These drugs block estrogen from connecting to the cancer cells and telling them to grow and divide. While they have anti-estrogen effects in breast cells, they act like an estrogen in other tissues, like the uterus and the bones.

These drugs are pills, taken by mouth.

Tamoxifen

Tamoxifen can be used to treat women with breast cancer who have or have not gone through menopause.

This drug can be used in several ways:

- In women at high risk of breast cancer, tamoxifen can be used to help lower the risk of developing breast cancer.
- For women who have been treated with breast-conserving surgery for ductal carcinoma in situ (DCIS) that is hormone receptor-positive, taking tamoxifen for 5 years lowers the chance of the DCIS coming back in the same breast. It also lowers the chance of getting an invasive breast cancer or another DCIS in both breasts.
- For women with hormone receptor-positive invasive breast cancer treated with surgery, tamoxifen can help lower the chances of the cancer coming back and improve the chances of living longer. It can also lower the risk of a new cancer developing in the other breast. Tamoxifen can be started either after surgery (adjuvant therapy) or before surgery (neoadjuvant therapy). When given after surgery, it is usually taken for 5 to 10 years. This drug is used mainly for women with early-stage breast cancer who have not yet gone through menopause. If you have gone through menopause, aromatase inhibitors (see below) are often used instead.
- For women with hormone-positive breast cancer that has spread to other parts of the body, tamoxifen can often help slow or stop the growth of the cancer, and might even shrink some tumors.

Toremifene (Fareston)

almost all women with hormone receptor-positive breast cancer.

Selective estrogen receptor degraders (SERDs)

Like SERMs, these drugs attach to estrogen receptors. But SERDs bind to the receptors more tightly and cause them to be broken down. These drugs have anti-estrogen effects throughout the body.

SERDs are used most often in women who are past menopause. When given to premenopausal women, they need to be combined with a luteinizing-hormone releasing hormone (LHRH) agonist to turn off the ovaries (see Ovarian suppression below).

Fulvestrant (Faslodex)

Fulvestrant can be used:

- Alone to treat advanced breast cancer that has not been treated with other hormone therapy.
- Alone to treat advanced breast cancer after other hormone drugs (like tamoxifen and often an aromatase inhibitor) have stopped working.
- In combination with a [CDK 4/6 inhibitor](#) or [PI3K inhibitor](#) to treat metastatic breast cancer as initial hormone therapy or after other hormone treatments have been tried.

It is given as 2 injections into the buttocks (bottom). For the first month, the 2 shots are given 2 weeks apart. After that, they are given once a month.

Elacestrant (Orserdu)

This drug can be used to treat advanced, ER-positive, HER2-negative breast cancer when the cancer cells have an [ESR1 gene mutation](#)⁴, and the cancer has grown after at least one other type of hormone therapy.

Elacestrant is taken daily as pills.

Side effects of fulvestrant and elacestrant

Common short-term side effects of these drugs can include:

- Hot flashes and/or night sweats
- Headache
- Nausea
- Feeling tired
- Loss of appetite
- Muscle, joint, or bone pain
- Injection site pain

Elacestrant can also increase cholesterol and fat levels in the blood.

Drugs that lower estrogen levels

Because estrogen stimulates hormone receptor-positive breast cancers to grow, lowering the estrogen level can help slow the cancer's growth or help prevent it from coming back.

Aromatase inhibitors (AIs)

Aromatase inhibitors (AIs) are drugs that stop most estrogen production in the body. Before menopause, most estrogen is made by the ovaries. But in women whose ovaries aren't working, either because they have gone through menopause or because of certain treatments, estrogen is still made in body fat by an enzyme called aromatase. AIs work by preventing aromatase from making estrogen.

These drugs are useful for women who have gone through menopause, although they can also be used in pre-menopausal women when they are combined with ovarian suppression (see below).

These AIs are pills taken every day to treat breast cancer:

- Letrozole (Femara)
- Anastrozole (Arimidex)
- Exemestane (Aromasin)

Possible side effects of AIs

The most common side effects of AIs are:

- Hot flashes
- Vaginal dryness
- Bone and joint pain
- Muscle pain

Als tend to have side effects different from tamoxifen. They don't cause uterine cancers and very rarely cause blood clots. They can, however, cause **muscle pain and joint stiffness and/or pain**

- **Chemotherapy drugs:** Some chemo drugs can damage the ovaries of premenopausal women so they no longer make estrogen. Ovarian function can return months or years later in some women, but in others the damage to the ovaries is permanent and leads to menopause.

All of these methods can cause symptoms of menopause, including hot flashes, night sweats, vaginal dryness, and mood swings.

Hormone therapy after surgery for breast cancer

After surgery for treatment of hormone receptor-positive breast cancer, hormone therapy can be given to reduce the risk of the cancer coming back.

These hormone therapy schedules are known to be helpful for women who are **postmenopausal when diagnosed**:

- An AI for 5 to 10 years
- An AI for 2 to 3 years, followed by tamoxifen for 2 to 3 years (5 years total of treatment)
- Tamoxifen for 2 to 3 years, followed by an AI for 2 to 3 years (5 years total of treatment)
- Tamoxifen for 2 to 3 years, followed by an AI for 5 years (7 to 8 years of treatment)
- Tamoxifen for 4½ to 6 years, followed by an AI for 5 years (9½ to 11 years of treatment)
- Tamoxifen for 5 to 10 years
- For women who are unable to take an AI, tamoxifen for 5 to 10 years is an option
- An AI along with ribociclib (Kisqali) for 3 years followed by AI alone to complete 5 years.

For most postmenopausal women whose cancers are hormone receptor-positive, most doctors recommend taking an AI at some point during adjuvant (after surgery) therapy. Standard treatment is to take these drugs for about 5 years, or to take in sequence with tamoxifen for 5 to 10 years. For women at a higher risk of recurrence, hormone treatment for longer than 5 years may be recommended. Tamoxifen is an option for some women who cannot take an AI. Taking tamoxifen for 10 years is considered more effective than taking it for 5 years, but you and your doctor will decide the best schedule of treatment for you.

These therapy schedules are known to be helpful for women who are **pre-menopausal when diagnosed**:

- Tamoxifen (with or without ovarian suppression) for 5 to 10 years.

Therapy⁵.

To learn about some of the side effects listed here and how to manage them, see [Managing Cancer-related Side Effects](#)⁶.

Hyperlinks

1. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-gene-expression.html
2. www.cancer.org/cancer/types/endometrial-cancer.html
3. www.cancer.org/cancer/types/uterine-sarcoma.html
4. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/other-breast-cancer-gene-protein-blood-tests.html
5. www.cancer.org/cancer/managing-cancer/treatment-types/hormone-therapy.html
6. www.cancer.org/cancer/managing-cancer/side-effects.html

References

Burstein HJ, Lacchetti C, Anderson H, Buchholz TA, Davidson NE, Gelmon KA et al. Adjuvant Endocrine Therapy for Women With Hormone Receptor-Positive Breast Cancer: ASCO Clinical Practice Guideline Focused Update. *J Clin Oncol*. 2019 Feb 10;37(5):423-438.

Burstein HJ, Lacchetti C, Anderson H, Buchholz TA, Davidson NE, Gelmon KA et al. Adjuvant Endocrine Therapy for Women With Hormone Receptor-Positive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update on Ovarian Suppression. *J Clin Oncol*. 2016;34(14):1689-1701.

to 10 years versus stopping at 5 years in 6,953 women with early breast cancer. *J Clin Oncol* (Meeting Abstracts) June 2013 vol. 31 no. 18_suppl 5.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Ma CX and Sparano JA. Treatment approach to metastatic hormone receptor-positive, HER2-negative breast cancer: Endocrine therapy and targeted agents. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Accessed August 11, 2021.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Health Professional Version. 2021. Accessed at <https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on August 11, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 6.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 11, 2021.

Osborne CK. Chapter 53: Adjuvant Systemic Therapy Treatment Guidelines. In: Harris JR, Lippman ME, Morrow M, Osborne CK, eds. *Diseases of the Breast*. 5th ed. Philadelphia: Wolters Kluwer Health; 2014.

Rimawi MF and Osborne CK. Chapter 43: Adjuvant Systemic Therapy: Endocrine Therapy. In: Harris JR, Lippman ME, Morrow M, Osborne CK, eds. *Diseases of the Breast*. 5th ed. Philadelphia: Wolters Kluwer Health; 2014.


Rocca A, Maltoni R, Bravaccini S, Donati C, Andreis D. Clinical utility of fulvestrant in the treatment of breast cancer: a report on the emerging clinical evidence. *Cancer Manag Res*. 2018;10:3083–3099.

Stearns V and Davidson NE. Chapter 45: Adjuvant Chemo Endocrine Therapy. In: Harris JR, Lippman ME, Morrow M, Osborne CK, eds. *Diseases of the Breast*. 5th ed. Philadelphia: Wolters Kluwer Health; 2014.

Last Revised: September 22, 2024

Targeted Drug Therapy for Breast Cancer

Targeted drug therapy uses medicines that are directed at (target) proteins on breast cancer cells that help them grow, spread, and live longer. Targeted drugs work to destroy cancer cells or slow down their growth. They have side effects different from [chemotherapy](#).



can't be removed with surgery or that has spread to another part of the body, typically after chemotherapy has been tried or if the cancer recurs within 6 months of finishing adjuvant chemotherapy.

Kinase inhibitors

HER2 is a type of protein known as a kinase. Kinases are proteins in cells that normally relay signals (such as telling the cell to grow). Drugs that block kinases are called **kinase inhibitors**.

Lapatinib (Tykerb): This drug is a pill taken daily. Lapatinib is used to treat advanced

Because these drugs can cause heart damage, doctors often check your heart function (with an echocardiogram or a MUGA scan) before treatment, and regularly while you are taking the drug. Let your doctor know if you develop symptoms such as **shortness**

hormone therapy drug.

Everolimus is taken as a pill, typically once a day.

Common **side effects of everolimus** include mouth sores, rash, diarrhea, nausea, feeling weak or tired, low blood counts, shortness of breath, and cough. Everolimus can also increase blood lipids (cholesterol and triglycerides) and blood sugars, so your doctor will check your blood work regularly while you are taking this drug. It can also increase your risk of serious infections, so your doctor will watch you closely for infection.

PI3K inhibitors

Alpelisib (Piqray) and **inavolisib (Itovebi)** block a form of the PI3K protein in cancer cells. This can help stop them from growing.

These drugs can be used to treat women with advanced hormone receptor-positive, HER2-negative breast cancer with a *PIK3CA* gene mutation that has grown during or after treatment with hormone therapy. (The *PIK3CA* gene is the gene that tells the cell to make the PI3K protein.) Your doctor will test your blood or tumor for this mutation before starting treatment with one of these drugs.

- Alpelisib is used along with the hormone drug fulvestrant.
- Inavolisib is used along with fulvestrant and the targeted drug palbociclib (see above).

These drugs are taken as pills, typically once a day.

Side effects of PI3K inhibitors can include skin rash, mouth sores, high blood sugar levels, signs of kidney, liver, or pancreatic problems, diarrhea, low blood counts, nausea and vomiting, fatigue, decreased appetite, weight loss, low calcium levels, blood clotting problems, and hair loss.

Some side effects can be more severe, and they might be more likely with one drug than with another. For example:

- Alpelisib is more likely to cause severe skin reactions, such as rashes with peeling and blistering, as well as serious lung inflammation (pneumonitis).
- Inavolisib is more likely to cause severe mouth sores.
- Both drugs might cause a person to have very high blood sugar levels or severe

diarrhea.

diarrhea.

This ADC can be used to treat advanced hormone receptor-positive, HER2-negative breast cancer, in people who have already received hormone therapy and at least 2 chemo regimens.

This drug is given into a vein (IV) weekly for 2 weeks, followed by one week off, then restarted.

Some common **side effects** of this drug include nausea, vomiting, diarrhea, constipation, feeling tired, rash, loss of appetite, hair loss, low red blood cell counts, and belly pain. Serious side effects can include very low white blood cell counts (with an increased risk of infection), severe diarrhea, and infusion reactions (similar to an allergic reaction) when the drug is infused. Medicines are normally given before each treatment to lower the chances of vomiting and infusion reactions.

Targeted therapy for women with *BRCA* gene mutations

Olaparib (Lynparza) and **talazoparib (Talzenna)** are drugs known as *PARP inhibitors*. PARP proteins normally help repair damaged [DNA inside cells](#)³. The *BRCA* genes (*BRCA1* and *BRCA2*) also help repair DNA (in a slightly different way), but mutations in one of those genes can stop this from happening. PARP inhibitors work by blocking the PARP proteins. Because tumor cells with a mutated *BRCA* gene already have trouble repairing damaged DNA, blocking the PARP proteins often leads to the death of these cells. These drugs are pills and are taken once or twice a day. They can be used in different ways to treat breast cancer.

- Olaparib can be given to women with a *BRCA* mutation with early-stage HER2-negative breast cancer after surgery who have been treated with chemotherapy (before or after surgery) and are at high risk of the cancer recurring. It is typically given for one year. When given in this way, it can help some women live longer.
- Olaparib and talazoparib can be used to treat advanced or metastatic, HER2-negative breast cancer in women with a *BRCA* mutation who have already had chemotherapy. If the cancer is hormone receptor-positive, olaparib can also be used in women who have already received hormone therapy.

Only a small portion of women with breast cancer are born with a [mutated *BRCA* gene](#),⁴ which is in all the cells of the body. This is different from the gene change happening after you are born which is found only in the cancer cells. If you are not known to have a *BRCA* mutation, your doctor will test your blood to be sure you have one before starting treatment with these drugs.

Side effects can include nausea, vomiting, diarrhea, fatigue, loss of appetite, taste changes, low red blood cell counts (anemia), low platelet counts, and low white blood cell counts. Rarely, some people treated with a PARP inhibitor have developed a blood cancer, such as [myelodysplastic syndrome](#)⁵ or [acute myeloid leukemia \(AML\)](#)⁶.

Targeted therapy for triple-negative breast cancer

In triple-negative breast cancer (TNBC), the cancer cells don't have estrogen or progesterone receptors and they make very little or none of the HER2 protein.

Antibody-drug conjugate

An antibody-drug conjugate (ADC) is a monoclonal antibody joined to a chemotherapy drug. The antibody acts like a homing signal by attaching to a specific protein on cancer cells, bringing the chemo directly to them.

Sacituzumab govitecan (Trodelvy): In the case of this ADC, the monoclonal antibody part attaches to the Trop-2 protein on breast cancer cells and brings a chemo drug, similar to irinotecan, directly to them. (Some breast cancer cells have too much Trop-2, which helps them grow and spread quickly.)

This antibody-drug conjugate can be used by itself to treat advanced TNBC, after at least 2 other chemo regimens have been tried. This drug is given in a vein (IV) weekly for 2 weeks, followed by one week off, then restarted.

Some common **side effects** of this drug include nausea, vomiting, diarrhea, constipation, feeling tired, rash, loss of appetite, hair loss, low red blood cell counts, and belly pain. Serious side effects can include very low white blood cell counts (with an increased risk of infection), severe diarrhea, and infusion reactions (similar to an allergic reaction) when the drug is infused. Medicines are normally given before each treatment to lower the chances of vomiting and infusion reactions.

More information about targeted therapy

To learn more about how targeted drugs are used to treat cancer, see [Targeted Cancer Therapy](#)⁷.

To learn about some of the side effects listed here and how to manage them, see [Managing Cancer-related Side Effects](#)⁸.

Hyperlinks

1. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-her2-status.html
2. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-her2-status.html
3. www.cancer.org/cancer/managing-cancer/treatment-types/biosimilar-drugs/list.html
4. www.cancer.org/cancer/understanding-cancer/genes-and-cancer.html
5. www.cancer.org/cancer/types/breast-cancer/risk-and-prevention/breast-cancer-risk-factors-you-cannot-change.html
6. www.cancer.org/cancer/types/myelodysplastic-syndrome.html
7. www.cancer.org/cancer/types/acute-myeloid-leukemia.html
8. www.cancer.org/cancer/managing-cancer/treatment-types/targeted-therapy.html
9. www.cancer.org/cancer/managing-cancer/side-effects.html

References

Almuwaqqat Z, Meisel JL, Barac A, Parashar S. Breast Cancer and Heart Failure. *Heart Fail Clin*. 2019 Jan;15(1):65-75.

Bardia A, Mayer IA, DiRHD0 giu 0 6 et al. ETIIMMU-132) in.83 vily Prerapy.ednd titmls 0 0 0 rg /GS30

DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Ma CX and Sparano JA. Treatment approach to metastatic hormone receptor-positive, HER2-negative breast cancer: Endocrine therapy and targeted therapy. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated August 10, 2021. Accessed August 13, 2021.

Morgan JP. Cardiotoxicity of trastuzumab and other HER2-targeted agents. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Accessed August 12, 2021.

Mukohara T. PI3K mutations in breast cancer: prognostic and therapeutic implications. *Breast Cancer (Dove Med Press)*. 2015;7: 111–123.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Health Professional Version. 2021. Accessed at <https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on August 12, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 6.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 13, 2021.

Schott AF. Systemic treatment for HER2-positive metastatic breast cancer. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Accessed August 12, 2021.

Last Revised: October 11, 2024

Immunotherapy for Breast Cancer

Immunotherapy is the use of medicines to boost a person's own immune system to recognize and destroy cancer cells more effectively. Immunotherapy typically works on specific proteins involved in the immune system to enhance the immune response. These drugs have side effects different from those of chemotherapy.

Some immunotherapy drugs, for example, monoclonal antibodies, work in more than one way to control cancer cells and may also be considered [targeted therapy](#) because

they block a specific protein on the cancer cell to keep it from growing.

Immunotherapy is used to treat some types of breast cancer.

- [Immune checkpoint inhibitors for breast cancer](#)
- [More information about immunotherapy](#)

Immune checkpoint inhibitors for breast cancer

An important part of the immune system is its ability to keep itself from attacking normal cells in the body. To do this, it uses proteins (or "checkpoints") on immune cells that need to be turned on (or off) to start an immune response. Breast cancer cells sometimes use these checkpoints to avoid being attacked by the immune system. Drugs that target these checkpoint proteins, help restore the immune response against breast cancer cells.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 6.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 13, 2021.

Schmidt P, Adams S, Rugo HS, Scheeweiss A, Barrios CH, Iwata H, et al. Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer. *N Engl J Med*. 2018 Nov 29;379(22):2108-2121. doi: 10.1056/NEJMoa1809615. Epub 2018 Oct 20.

Last Revised: October 27, 2021

Treatment of Breast Cancer by Stage

This information is based on AJCC Staging systems prior to 2018 which were primarily based on tumor size and lymph node status. Since the updated staging system for breast cancer now also includes estrogen receptor (ER), progesterone receptor (PR), and HER2 status, the stages may be higher or lower than previous staging systems. Whether or not treatment strategies will change with this new staging system are yet to be determined. You should discuss your stage and treatment options with your doctor.

The [stage](#)¹ of your breast cancer is an important factor in making decisions about your treatment options. In general, the more the breast cancer has spread, the more treatment you will likely need. But other factors can also be important, such as:

- If the cancer cells have [hormone receptors](#)² (that is, if the cancer is ER-positive or PR-positive)
- If the cancer cells have large amounts of the [HER2 protein](#)³ (that is, if the cancer is HER2-positive)
- If the cancer cells have a certain gene mutation (change)
- Your overall health and personal preferences
- If you have gone through menopause or not

Stage 0 cancers are limited to the inside of the milk duct and are non-invasive (does not invade nearby tissues). Ductal carcinoma in situ (DCIS) is a stage 0 breast tumor. Lobular carcinoma in situ (LCIS) used to be categorized as stage 0, but this has been changed because it is not cancer. Still, it does indicate a higher risk of breast cancer. See Lobular Carcinoma in Situ (LCIS) for more information.

- [Treatment of Ductal Carcinoma in Situ \(DCIS\)](#)

[Stages I-III](#)

Treatment for stages I to III breast cancer usually includes surgery and radiation

Treatment of Ductal Carcinoma in Situ (DCIS)

Ductal carcinoma in situ (DCIS) means the cells that line the milk ducts of the breast have become cancer, but they have not spread into surrounding breast tissue.

DCIS is considered non-invasive or pre-invasive breast cancer. DCIS can't spread outside the breast, but it is often treated because if left alone, some DCIS cells can continue to undergo abnormal changes that cause it to become invasive breast cancer (which can spread).

In most cases, a woman with DCIS can choose between breast-conserving surgery (BCS) and simple mastectomy. But sometimes, if DCIS is throughout the breast, a mastectomy might be a better option. There are clinical studies being done to see if observation instead of surgery might be an option for some women.

- [Breast-conserving surgery \(BCS\)](#)
- [Mastectomy](#)
- [Hormone therapy after breast surgery](#)

Breast-conserving surgery (BCS)

In [breast-conserving surgery](#) [sentinel lymph node biopsy \(SLNB\)](#) (BCS), the surgeon removes the tumor and a small amount of normal breast tissue around it. Lymph node removal is not usually needed with BCS. It might be done after the first surgery if an

Mastectomy

Simple [mastectomy](#) (removal of the entire breast) may be needed if the area of DCIS is very large, if the breast has several separate areas of DCIS in different quadrants (multicentric), or if BCS cannot remove the DCIS completely (that is, the BCS specimen and re-excision specimens still have cancer cells in or near the surgical margins). If a mastectomy is needed for any of the reasons stated above, many doctors will do a SLNB along with the mastectomy because there is a higher chance that invasive cancer might be found. If an area of invasive cancer is found in the tissue removed during a mastectomy, the doctor won't be able to go back and do SLNB later, and as a result may have to do a full axillary lymph node dissection (ALND) instead.

Women having a mastectomy for DCIS typically don't need radiation therapy and may choose to have [breast reconstruction](#)¹ right away or later.

Hormone therapy after breast surgery

If the DCIS is hormone receptor-positive (estrogen or progesterone), treatment with tamoxifen (for any woman) or an aromatase inhibitor, such as exemestane or anastrozole, (for women past menopause) for 5 years after surgery can lower the risk of another DCIS or invasive cancer developing in either breast. If you have hormone receptor-positive DCIS, discuss the reasons for and against hormone therapy with your doctor.

Hyperlinks

1. www.cancer.org/cancer/types/breast-cancer/reconstruction-surgery.html

References

Collins LC, Laronga C, and Wong JS. Ductal carcinoma in situ: Treatment and prognosis. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated July 13, 2021. Accessed August 17, 2021.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant

Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*.

Treatment of Breast Cancer Stages I-III

- [Treating stage I breast cancer](#)
- [Treating stage II breast cancer](#)
- [Treating stage III breast cancer](#)

What type of drug treatment(s) might I get?

Most women with breast cancer in stages I, II, or III will get some kind of systemic therapy as part of their treatment. This might include:

- [Chemotherapy](#)
- [Hormone therapy](#) (tamoxifen, an aromatase inhibitor, or one followed by the other)
- [Targeted drugs](#), such as trastuzumab (Herceptin), pertuzumab (Perjeta), or abemaciclib (Verzenio)
- [Immunotherapy](#)
- Some combination of these

The types of drugs that might work best depend on the tumor's hormone receptor status, HER2 status, and other factors.

Treating stage I breast cancer

These breast cancers are still fairly small and either have not spread to the lymph nodes or have spread to only a tiny area in the sentinel lymph node (the first lymph node to which cancer is likely to spread).

Local therapy (surgery and radiation therapy)

Surgery is the main treatment for stage I breast cancer. These cancers can be treated with either [breast-conserving surgery](#) (BCS; sometimes called lumpectomy or partial mastectomy) or [mastectomy](#). The nearby [lymph nodes](#) will also need to be checked, either with a sentinel lymph node biopsy (SLNB) or an axillary lymph node dissection (ALND).

Some women can have [breast reconstruction](#)⁶ at the same time as the surgery to remove the cancer. But if you will need radiation therapy after surgery, it is better to wait to get reconstruction until after the radiation is complete.

If BCS is done, radiation therapy is usually given after surgery to lower the chance of the cancer coming back in the breast and to also help people live longer.

In a separate group, women who are at least 65 years old may consider BCS **without** radiation therapy if ALL of the following are true:

- The tumor was 3 cm (a little more than 1 inch) or less across and it has been removed completely.
- None of the lymph nodes removed contained cancer.
- The cancer is ER-positive or PR-positive, and hormone therapy will be given.

Radiation therapy given to women with these characteristics still lowers the chance of the cancer coming back, but it has not been shown to help them live longer.

If you had a mastectomy, you are less likely to need radiation therapy, but it might be given depending on the details of your specific cancer. You should discuss if you need radiation treatment with your doctor. You might be sent to a doctor who specializes in radiation (a radiation oncologist) for evaluation.

Systemic therapy (chemo and other drugs)

If a woman has a hormone receptor-positive (ER-positive or PR-positive) breast cancer, most doctors will recommend hormone therapy (tamoxifen or an aromatase inhibitor, or one followed by the other) as an adjuvant (after surgery) treatment, no matter how small the tumor is. Women with tumors larger than 0.5 cm (about ¼ inch) across may be more likely to benefit from it. Hormone therapy is typically given for at least 5 years.

If the tumor is larger than 0.5 cm (about 1/4 inch) across, chemo after surgery (adjuvant chemotherapy) is sometimes recommended. A woman's age when she is diagnosed may help in deciding if chemo should be offered or not. Some doctors may suggest chemo for smaller tumors as well, especially if they have any unfavorable features (a cancer that is growing fast; hormone receptor-negative, HER2-positive; or having a [high score on a gene panel such as Oncotype DX⁷](#)).

After surgery, some women with HER2-positive cancers will be treated with trastuzumab (with or without pertuzumab) for up to 1 year.

Many women with HER2-positive cancers will be treated with neoadjuvant (before surgery) chemo and trastuzumab (with or without pertuzumab) followed by surgery and more trastuzumab (with or without pertuzumab) for up to 1 year. If after neoadjuvant therapy, residual cancer is found during surgery, trastuzumab may be changed to a different drug, called ado-trastuzumab emtansine, which is given every 3 weeks for 14 doses.

For women with a BRCA mutation and hormone-positive, HER2-negative breast cancer who received neoadjuvant chemotherapy but still have residual cancer at the time of surgery, the targeted drug olaparib might be given after surgery. It is usually given for one year. When given this way, it can help some women live longer.

Treating stage II breast cancer

Stage II breast cancers are larger than stage I cancers and/or have spread to a few nearby lymph nodes.

Local therapy (surgery and radiation therapy)

Stage II cancers are treated with either breast-conserving surgery (BCS; sometimes called lumpectomy or partial mastectomy) or mastectomy. The nearby [lymph nodes](#) will also be checked, either with a sentinel lymph node biopsy (SLNB) or an axillary lymph node dissection (ALND).

Women who have [BCS](#) are treated with radiation therapy after surgery. Women whoALND).



at surgery after neoadjuvant chemo, the targeted drug olaparib might be given for one year to help lower the chance of the cancer recurring. When given this way, it can help some women live longer.

- **Immunotherapy:** Women with TNBC might get the immunotherapy drug, pembrolizumab, before surgery and then again after surgery. See [Treatment of Triple-negative Breast Cancer](#) for more details.

Treating stage III breast cancer

In stage III breast cancer, the tumor is large (more than 5 cm or about 2 inches across) or growing into nearby tissues (the skin over the breast or the muscle underneath), or the cancer has spread to many nearby lymph nodes.

If you have inflammatory breast cancer: Stage III cancers also include some inflammatory breast cancers that have not spread beyond nearby lymph nodes. These cancers are treated slightly different from other stage III breast cancers. You can find more details in [Treatment of Inflammatory Breast Cancer](#).

There are two main approaches to treating stage III breast cancer:

Starting with neoadjuvant therapy



ado-trastuzumab emtansine may be used instead of trastuzumab. It is given every 3 weeks for 14 doses. For women with hormone receptor-positive cancer that is in the lymph nodes, who have completed a year of trastuzumab, the doctor might also recommend additional treatment with an oral targeted drug called neratinib for a year.

Women with hormone receptor-positive (ER-positive or PR-positive) breast cancers will also get adjuvant hormone therapy which can typically be taken at the same time as trastuzumab.

For women with hormone receptor-positive, HER2-negative breast cancer that is in the lymph nodes, and has a high chance of coming back, abemaciclib can be given after surgery along with tamoxifen or an AI. It is a pill typically given twice a day for 2 years.

For women who have a BRCA mutation and hormone receptor-positive, HER2-negative breast cancer and still have cancer in the tissue removed at surgery after neoadjuvant chemo, the targeted drug olaparib might be given for one year to help lower the chance of the cancer recurring. When given this way, it can help some women live longer.

Neoadjuvant treatment is a preferable option for women with stage III TNBC or HER2-positive breast cancer because the treatment given after surgery is chosen depending on how much cancer is still in the breast and/or lymph nodes at the time of surgery. Some women with stage III cancer who get neoadjuvant treatment might live longer if the cancer goes away completely with that treatment.

Women with TNBC might get the immunotherapy drug, pembrolizumab, before surgery and then again after surgery. See [Treatment of Triple-negative Breast Cancer](#) for more details.

Starting with surgery

Surgery first is an option for some women with stage III cancers. Because these tumors are fairly large and/or have grown into nearby tissues, this usually means getting a mastectomy. For women with fairly large breasts, BCS may be an option if the cancer hasn't grown into nearby tissues. SLNB may be an option for some patients, but most will need an ALND. Surgery is usually followed by adjuvant chemotherapy, and/or hormone therapy, and/or targeted drug therapy, and/or HER2-positive treatment (trastuzumab, pertuzumab, or neratinib) depending on the traits of the cancer cells. Radiation is recommended after surgery.

Hyperlinks

1. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/stages-of-breast-cancer.html
2. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-hormone-receptor-status.html
3. www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-her2-status.html

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 7.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 27, 2021.

Sparano JA, Gray RJ, Makower KI, Pritchard KS, Albain DF, Hayes CE, et al. Adjuvant chemotherapy guided by a 21-gene expression assay in breast cancer [published online ahead of print June 3 2018]. *NEJM*. 2018; doi:

Treatment of Stage IV (Metastatic) Breast Cancer

- [Systemic \(drug\) treatments for stage IV breast cancer](#)
- [Local or regional treatments for stage IV breast cancer](#)
- [Relieving symptoms of advanced breast cancer](#)
- [Advanced cancer that progresses during treatment](#)

Systemic (drug) treatments for stage IV breast cancer

Treatment often continues until the cancer starts growing again or until side effects become unacceptable. If this happens, other drugs might be tried. The types of drugs used for stage IV breast cancer depend on the hormone receptor status, the HER2 status of the cancer, and sometimes gene mutations that might be found.

Hormone receptor-positive cancers

Women with hormone (estrogen or progesterone) receptor-positive cancers are sometimes treated first with hormone therapy (tamoxifen or an aromatase inhibitor). This may be combined with a targeted drug such as a CDK4/6 inhibitor.

If the initial hormone therapy stops working, another type can be tried, often along with a targeted drug such as everolimus, a PI3K inhibitor such as alpelisib (Piqray) or inavolisib (Itovebi), or an AKT inhibitor such as capivasertib (Truqap), if the cancer cells have certain gene changes. (See “Advanced cancer that progresses during treatment” below.)

Women who haven’t yet gone through menopause are often treated with tamoxifen or with medicines that keep the ovaries from making hormones along with other drugs.

Hormone receptor-negative cancers

Chemo is the main treatment for women with hormone (estrogen and progesterone) receptor-negative cancers, because hormone therapy isn’t helpful for these cancers.

HER2-positive cancers

The first therapy given is usually chemotherapy in combination with trastuzumab (Herceptin, [other names](#)¹) and pertuzumab (Perjeta), both HER2 targeted drugs. If the cancer grows, other options might include:

- An [antibody-drug conjugate](#)
- A [kinase inhibitor](#) with an anti-HER2 drug or with a [chemo](#) drug or both

For hormone receptor-positive (ER-positive or PR-positive) cancers that were being treated with hormone therapy, switching to another type of hormone therapy and/or adding a [targeted drug](#) sometimes helps. For example:

- If either letrozole (Femara) or anastrozole (Arimidex) was given first, using exemestane, possibly with the targeted drug everolimus (Afinitor), may be an option.
- Another option might be using elacestrant (Orserdu), fulvestrant (Faslodex), or a different aromatase inhibitor, sometimes along with a targeted drug known as a CDK4/6 inhibitor.
- If the cancer has a *PIK3CA* gene mutation and has grown while being treated with an aromatase inhibitor, fulvestrant with the targeted drug alpelisib (Piqray) might be considered. Another option might be the combination of fulvestrant with the targeted drugs palbociclib (Ibrance) and inavolisib (Itovebi).
- If the cancer has a *PIK3CA*, *AKT1*, or *PTEN* gene mutation, fulvestrant with the targeted drug capivasertib (Truqap) might be an option.

If the cancer is no longer responding to any hormone drugs, chemotherapy immunotherapy, or PARP inhibitors might be options depending on specific features of the cancer or any gene changes that might be present.

Progression during chemotherapy

If the cancer is no longer responding to one chemo regimen, trying another may be helpful. Many different drugs and combinations can be used to treat breast cancer. However, each time a cancer progresses during treatment, it becomes less likely that further treatment will have an effect. Sometimes, other options include adding an immunotherapy drug to the chemo or using a PARP inhibitor alone depending on specific features of the cancer or any gene changes that might be present.

Progression while being treated with HER2 drugs

HER2-positive cancers that no longer respond to trastuzumab (Herceptin) might respond to other drugs that target the HER2 protein. Options for women with HER2-positive cancers might include:

- Margetuximab (Margenza) with chemo
- Lapatinib (Tykerb) and the oral chemo drug capecitabine
- Lapatinib and an aromatase inhibitor (for hormone receptor-positive cancers)
- Neratinib (Nerlynx) and the chemo drug capecitabine (this combination can be helpful for cancers that have spread to the brain)
- Tucatinib (Tukysa), trastuzumab, and the chemo drug capecitabine (this combination can be helpful for cancers that have spread to the brain)

Because current treatments are very unlikely to cure metastatic breast cancer, if you are in otherwise good health, you may want to think about taking part in a [clinical trial](#)⁴ testing a newer treatment.

Hyperlinks

1. www.cancer.org/cancer/managing-cancer/treatment-types/biosimilar-drugs/list.html
www.cancer.org/cancer/types/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-her2-status.html

<https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on August 30, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 7.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 30, 2021.

Last Revised: October 11, 2024

Treatment of Recurrent Breast Cancer

For some women, breast cancer may come back after treatment – sometimes years later. This is called a recurrence. **Recurrence can be local (in the same breast or in the surgery scar), regional (in nearby lymph nodes), or in a distant area.** Cancer that is found in the opposite breast without any cancer elsewhere in the body is not a recurrence—it is a new cancer that requires its own treatment.

- [Treating local recurrence](#)
- [Treating regional recurrence](#)
- [Treating distant recurrence](#)

Treating local recurrence

For women whose breast cancer has recurred locally, treatment depends on their initial treatment.

- If you had breast-conserving surgery (lumpectomy), a local recurrence in the breast is usually treated with [mastectomy](#).
- If the initial treatment was mastectomy, recurrence near the mastectomy site is treated by removing the tumor whenever possible. This is often followed by [radiation therapy](#) if not given before.

In either case, [hormone therapy](#), [targeted therapy](#) (like trastuzumab), [immunotherapy](#), [chemotherapy](#), or some combination of these may be used after surgery and/or radiation therapy. These drugs might also be used if surgery or radiation are not options.

Treating regional recurrence

When breast cancer comes back in nearby lymph nodes (such as those under the arm or around the collar bone), it is treated by [removing those lymph nodes](#), if possible. This may be followed by radiation aimed at the area if it was not given before. Systemic treatment (such as chemo, targeted therapy, or hormone therapy) may be considered after surgery as well.

Treating distant recurrence

In general, women whose breast cancer comes back in other parts of the body, such as the bones, lungs, or brain, are treated the same way as those found to have stage IV breast cancer in these organs when they were first diagnosed. See [Treating Stage IV \(Metastatic\) Breast Cancer](#). The only difference is that treatment may or may not include treatments or drugs a woman has already had.

Recurrent breast cancer can sometimes be hard to treat. If you are in otherwise good health, you might want to think about taking part in a [clinical trial](#)¹ testing a newer treatment.

Should your cancer come back, see [Understanding Recurrence](#)² for more on how to manage and cope with this phase of your treatment.

Hyperlinks

1. www.cancer.org/cancer/managing-cancer/making-treatment-decisions/clinical-trials.html
2. www.cancer.org/cancer/survivorship/long-term-health-concerns/recurrence.html

References

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Health Professional Version. 2021. Accessed at <https://www.cancer.gov/types/breast/hp/breast-treatment-pdq> on August 30, 2021.

National Comprehensive Cancer Network (NCCN). Practice Guidelines in Oncology: Breast Cancer. Version 7.2021. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on August 30, 2021.

Last Revised: October 27, 2021

Treatment of Triple-negative Breast Cancer

Triple-negative breast cancer (TNBC) doesn't have estrogen or progesterone receptors and also makes too little or none of the HER2 protein. Because the cancer cells don't have these proteins, hormone therapy and drugs that target HER2 are not helpful, so chemotherapy (chemo) is the main **systemic** treatment option. And even though TNBC tends to respond well to chemo initially, it often comes back or spreads to other parts of the body.

the [targeted drug olaparib \(Lynparza\)](#) might be given for a year after adjuvant chemo. When given this way, it can help some women live longer.

(changes) inside the cancer cells. Cells that have many gene mutations (a high TMB) might be more likely to be recognized as abnormal and attacked by the body's immune system.

[Surgery](#) and [radiation](#) may also be options in certain situations.

See [Treatment of Stage IV \(Metastatic\) Breast Cancer](#) for more information.

Recurrent triple-negative breast cancer

If TNBC comes back (recurs) locally, cannot be removed with surgery, and makes the PD-L1 protein, [immunotherapy](#) with the drug pembrolizumab along with chemotherapy is an option. Other treatments might be options as well, depending on the situation.

If the cancer recurs in other parts of the body, options might include chemotherapy or the [antibody-drug conjugate](#) sacituzumab govitecan (Trodelvy).

Regardless of the stage of the cancer, participation in a [clinical trial](#)¹ of new treatments for TNBC is also a good option because TNBC is uncommon and tends to have a poor prognosis (outcome) compared to other types of breast cancer, and because these studies often allow patients to have access to drugs not available for standard treatment.

Hyperlinks

1. www.cancer.org/cancer/managing-cancer/making-treatment-decisions/clinical-trials.html

References

Anders CK and Carey LA. ER/PR negative, HER2-negative (triple-negative) breast cancer. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated July 21, 2021. Accessed August 24, 2021.

Bardia A, Mayer IA, Diamond JR, et al. Efficacy and Safety of Anti-Trop-2 Antibody Drug Conjugate Sacituzumab Govitecan (IMMU-132) in Heavily Pretreated Patients With Metastatic Triple-Negative Breast Cancer. *J Clin Oncol*. 2017;35(19):21412148. doi:10.1200/JCO.2016.70.8297.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: *DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Jhan JR, Andrechek ER. Triple-negative breast cancer and the potential for targeted therapy. *Pharmacogenomics*. 2017;18(17):1595–1609.

Li X, Yang J, Peng L, Sahin AA, Huo L, Ward KC, O'Regan R, Torres MA, Meisel JL. Triple-negative breast cancer has worse overall survival and cause-specific survival than non-triple-negative breast cancer. *Breast Cancer Res Treat*. 2017 Jan;161(2):279-287.

National Cancer Institute. Physician Data Query (PDQ). Breast Cancer Treatment – Health Professional Version. 2021. Accessed at Breast Cancer .Version. .2321. Accessed at

Treatment of Inflammatory Breast Cancer

Inflammatory breast cancer (IBC) is an uncommon type of invasive breast cancer that typically makes the skin on the breast look red and feel warm. It also may give the breast skin a thick, pitted appearance that looks a lot like an orange peel. These changes are caused by cancer cells blocking lymph vessels in the skin.

Because inflammatory breast cancer has reached these lymph vessels and has caused changes in the skin, it is considered to be at least a stage III breast cancer when it is diagnosed. IBC that has spread to other parts of the body is considered stage IV. These

HER2), the [targeted therapy drug](#) trastuzumab (Herceptin) is usually given, sometimes along with another targeted drug, pertuzumab (Perjeta). These drugs can lead to heart problems when given with an anthracycline, so one option is to give the anthracycline first (without trastuzumab or pertuzumab), followed by treatment with a taxane and trastuzumab (with or without pertuzumab).

For IBC that is [triple-negative²](#), the [immunotherapy drug](#) pembrolizumab can be given with chemo before surgery (neoadjuvant treatment) and then continued by itself after surgery (adjuvant treatment).

For women who have:

- a BRCA mutation AND
- triple-negative or HER2-negative IBC AND
- residual cancer in the tissue removed by surgery after neoadjuvant chemo

the targeted drug olaparib (Lynparza) might be given to lower the risk of the cancer recurring. It is typically given for one year. When given this way, olaparib can help some women live longer.

Surgery and further treatments

If the cancer improves with chemo, [surgery](#) is typically the next step. The standard operation is a modified radical mastectomy, where the entire breast and the lymph nodes under the arm are removed. Because IBC affects so much of the breast and skin, breast-conserving surgery (partial mastectomy or lumpectomy) and skin-sparing mastectomy are not options. It isn't clear that sentinel lymph node biopsy (where only one or a few nodes are removed) is reliable in IBC, so it is also not an option.

If the cancer does not respond to chemo (and the breast is still very swollen and red), surgery cannot be done. Either other chemo drugs will be tried, or the breast may be treated with radiation. Then if the cancer responds (the breast shrinks and is no longer red), surgery may be an option.

If breast radiation isn't given before surgery, it is given after surgery, even if no cancer is thought to remain. This is called **adjuvant** radiation. It lowers the chance that the cancer will come back. Radiation is usually given 5 days a week for 6 weeks, but in some cases a more intense treatment (twice a day) can be used instead. Depending on how much tumor was found in the breast after surgery, radiation might be delayed until further chemo and/or targeted therapy (such as trastuzumab) is given. If breast reconstruction is to be done, it is usually delayed until after the radiation therapy that

Curigliano G. Inflammatory breast cancer and chest wall disease: The oncologist perspective. *Eur J Surg Oncol*. 2018 Aug;44(8):1142-1147.

Dawood S, Merajver SD, Viens P, et al. International expert panel on inflammatory breast cancer: Consensus statement for standardized diagnosis and treatment. *Ann Oncol*. 2011;22:515523.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds.

Treating Breast Cancer During Pregnancy

If you are diagnosed with breast cancer while pregnant, your treatment options will be more complicated because you will want to get the best treatment for your cancer while also protecting the baby. The type and timing of treatment will need to be planned carefully and coordinated between your cancer care team and your obstetrician.

When treating a pregnant woman with breast cancer, the goal is the same as when treating a non-pregnant woman: to cure the cancer whenever possible, or to control it and keep it from spreading if it can't be cured. But the extra concern of protecting a growing fetus may make treatment more complicated.

- [Is it safe to have breast cancer treatment during pregnancy?](#)
- [Breast cancer surgery during pregnancy](#)
- [Treatment after surgery](#)
- [Can I breastfeed during cancer treatment?](#)
- [Does pregnancy affect survival rates for breast cancer?](#)

Is it safe to have breast cancer treatment during pregnancy?

If you are pregnant and have breast cancer, you might have hard choices to make, so get expert help and be sure you know all your options. Pregnant women can safely get treatment for breast cancer, although the types of treatment used and the timing of treatment might be affected by the pregnancy. If you are pregnant and have been diagnosed with breast cancer, your treatment recommendations will depend on:

- The size of the tumor
- Where the tumor is located
- If the cancer has spread and if so, how far
- How far along you are in the pregnancy
- Your overall health
- Your personal preferences

It is generally safe to have surgery for breast cancer while you're pregnant. Chemotherapy seems to be safe for the baby if given in the second or third trimester of pregnancy, but it isn't safe in the first trimester. Other [breast cancer treatments](#), such as hormone therapy, targeted therapy, and radiation therapy, are more likely to harm the

these concerns, most experts recommend that SLNB only be used in certain situations such as later in pregnancy, and that the blue dye not be used during the procedure.

Is anesthesia safe during pregnancy?

Surgery for breast cancer generally carries little risk to the baby. But there are certain times in pregnancy when anesthesia (the drugs used to make you sleep for surgery) may be risky for the baby.

Your surgeon and anesthesiologist, along with a high-risk obstetrician (OB), will need to work together to decide the best time during pregnancy to operate. If the surgery is done later in the pregnancy, your obstetrician may be there just in case there are any problems with the baby during surgery. Together, your doctors will decide which anesthesia drugs and techniques are the safest for both you and the baby.

Treatment after surgery

Depending on the cancer's [stage](#)², you may need more treatment such as chemotherapy, radiation therapy, hormone therapy, and/or targeted therapy after surgery to help lower the risk of the [cancer coming back](#)³. This is called **adjuvant treatment**. In some cases, this treatment can be put off until after delivery.

Chemotherapy

Chemotherapy (chemo) may be used after surgery (as adjuvant treatment) for some earlier stages of breast cancer. It also can be used by itself for more advanced cancers.

Chemo is not given during the first 3 months (first trimester) of pregnancy. Because a lot of the baby's development occurs during this time, the safety of chemo hasn't been studied in the first trimester. The risk of miscarriage (losing the baby) is also the greatest during this time.

For many years, it was thought that all chemo would harm an unborn baby no matter when it was given. But studies have shown that certain [chemo](#) [chemo](#) [Tm](#) /[F2](#)(you may need more treat

will usually be delayed until at least your second trimester. If you are already in the third trimester when the cancer is found, chemo may be delayed until after birth. The birth may be induced (brought on) a few weeks early in some women. These same treatment plans may also be used for women with more advanced cancer.

Chemo is generally not recommended after 35 weeks of pregnancy or within 3 weeks of delivery because it can [lower the mother's blood cell counts](#)⁴. This could cause bleeding and increase the chances of infection during birth. Holding off on chemo for the last few weeks before delivery allows the mother's blood counts to return to normal before childbirth.

Treatments that typically are on hold until after delivery

Some treatments for breast cancer can harm the baby and are not safe during pregnancy. If these treatments are needed, they are usually scheduled after the baby is born.

Radiation therapy: Radiation therapy to the breast is often used after breast-conserving surgery (lumpectomy) to help reduce the risk of the cancer coming back. The high doses of radiation used for this can harm the baby any time during pregnancy. This may cause miscarriage, birth defects, slow fetal growth, or a higher risk of childhood cancer. Because of this, doctors don't use radiation treatment during pregnancy.

For some women whose cancer is found later in the pregnancy, it may be possible to have a lumpectomy during pregnancy and then wait until after the baby is born to get radiation therapy. But this treatment approach has not been well-studied. Waiting too long to start radiation can increase the chance of the cancer coming back.

Hormone therapy: Hormone therapy is often used as treatment after surgery or as treatment for advanced breast cancer in women with hormone receptor-positive (estrogen or progesterone) breast cancer. Hormone therapy drugs used for breast cancer include tamoxifen, anastrozole, letrozole, and exemestane.

Hormone therapy should not be given during pregnancy because it can affect the baby. It should be delayed until after the woman has given birth.

Targeted therapy: Drugs that target HER2, such as trastuzumab (Herceptin), pertuzumab (Perjeta), ado-trastuzumab emtansine (Kadcyla) and lapatinib (Tykerb), are important in treating HER2-positive breast cancers. In women who aren't pregnant, trastuzumab is used as a part of treatment after surgery, pertuzumab can be used with trastuzumab before surgery, and all of these drugs can be useful in treating advanced

cancer. But based on studies of women who were treated during pregnancy, none of these drugs are considered safe for the baby if taken during pregnancy.

Everolimus (Afinitor) and palbociclib (Ibrance) are also targeted drugs that can be used with hormone therapy to treat advanced breast cancer. Again, these and other targeted drugs are thought to be unsafe to use during pregnancy.

Can I breastfeed during cancer treatment?

Most doctors recommend that women who have just had babies and are about to be treated for breast cancer should stop (or not start) breastfeeding. Many chemo, hormone, and targeted therapy drugs can enter breast milk and be passed on to the baby. Breastfeeding isn't recommended if you are being treated with systemic drugs and sometimes shouldn't be restarted for months after treatment has ended. Sometimes, if a woman has hormone-positive breast cancer, she might be given drugs to stop the production of breast milk.

If breast surgery is planned, stopping breastfeeding will help reduce blood flow to the breasts and make them smaller. This can help with the operation. It also helps reduce the risk of infection in the breast and can help avoid breast milk collecting in biopsy or surgery areas.

If you have questions, such as when it might be safe to start breastfeeding, talk with your health care team. If you plan to start breastfeeding after you've stopped for a while, plan ahead. Breastfeeding (le7Dabies and argnan.n twsv8a.e team0 g dtll 7le7Da3ou h 0 0 rg td. ori6

be a difficult area to study. Finally, there are no reports showing that breast cancer itself can harm the baby.

Hyperlinks

Guidroz JA, Scott-Conner CEH, Weigel RJ. Management of pregnant women with breast cancer. *J Surg Oncol*. 2011;103:337-340.

Harlow SP and Weaver DL. Overview of sentinel lymph node biopsy in breast cancer. In Chen W, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated October 06, 2020. Accessed August 24, 2021.

Henry NL, Shah PD, Haider I, Freer PE, Jagsi R, Sabel MS. Chapter 88: Cancer of the Breast. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Jagsi R, King TA, Lehman C, Morrow M, Harris JR, Burstein HJ. Chapter 79: Malignant Tumors of the Breast. In: DeVita VT, Lawrence TS, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Johansson ALV, Andersson TM, Hsieh CC, Jirstrom K, Cnattingius S, Fredriksson I, et al. Tumor characteristics and prognosis in women with pregnancy-associated breast cancer. *Int J Cancer*. 2018 Apr 1;142(7):1343-1354.

Litton JK. Gestational breast cancer: Treatment. In Vora SR, ed. *UpToDate*. Waltham, Mass.: UpToDate, 2021. <https://www.uptodate.com>. Last updated November 19, 2020. Accessed August 24, 2021.

Loibl S, Han SN, von Minckwitz G, et al. Treatment of breast cancer during pregnancy: An observational study. *Lancet Oncol*. 2012;13:887-896.

McGrath SE, Ring A. Chemotherapy for breast cancer in pregnancy: evidence and guidance for oncologists. *Ther Adv Med Oncol*. 2011;3(2):73-83.

Hay (breast cancer.)Tj 0 g /F3 12 Tf 0 0 0 rg /G61606 gsJ Cancer. 201218:2 34-2 3019.

cancer.org | 1.800.227.2345